



The Important of Beta Carotene on Poultry Nutrition

Süleyman ÇALIŞLAR^{1*}

¹Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Animal Science, Kahramanmaraş, Turkey

ARTICLE INFO

Article history:

Received date: 27.05.2019

Accepted date: 08.08.2019

Edited by:

İbrahim AYTEKİN; *Selçuk University, Turkey*

Reviewed by:

Yusuf CUFADAR; *Selçuk University, Turkey*

Ahmet Engin TÜZÜN; *Adnan Menderes University, Turkey*

Keywords:

Beta carotene
Carotenoid
Poultry
Pigmentation
Antioxidant
Immunity

ABSTRACT

Beta carotene, the primary source of vitamin A in poultry rations, is one of the most important carotenoids. Under the influence of enzymes, Beta carotene (BC) is converted to vitamin A. The BC molecule is a double retinal structure and theoretically gives 2 molecules retinal. Its biological activity is only half of retinal. Conversion of carotenoids to retinol is rarely 100%. Thus the vitamins of various foods are expressed in terms of the potential retinol equivalence (RE).

BC is absorbed from the duodenum and if there is oil in the intestinal tract, it is absorbed faster. Oxidatively converting BC into vitamin A is mainly carried out in the intestinal brush border membrane, organs such as the liver, kidney and lungs. BC egg yolk is transported to and stored in immune organs and similar tissues. The BC content of the egg of the poultry varies. BC contents of hen eggs are low, while BC contents of eggs of wild birds are between 25-30%. Despite depletion of BC in the liver it's transfer to the egg continues.

Since poultry can not synthesize BC, it must be taken from outside. Products such as yellow corn, marigold and alfalfa are very rich sources of beta-carotene. BC is abundant in egg yolks.

BC is effective in the pigmentation of skin and egg yolks of hens. Due to BC's antioxidant properties prevents deterioration of egg and meat. It has also been shown that BC has important effects on the immunity and endocrine system. BC, strengthens see function, reduces the risk of cardiovascular disease, prevents inflammation and some types of cancer. Studies have shown that BC enhances the immune system by raising antibody response in poultries and prevents acute respiratory tract infections.

In this review article, the introduction of BC, its functions, effects on poultry nutrition were investigated.

1. Introduction

It has been observed that the nutrient profile (fatty acids, minerals, vitamins, etc.) of the egg, which is considered to be a highly nutritious food among nutritional sources, can be significantly improved by diet manipulation. For this reason, intensive research is carried out on the passage of some nutrients that can affect human health positively to eggs (Bean and Leeson, 2003; Khan et al., 2012).

It is stated that the enrichment of egg in terms of carotenoids will be beneficial for human health (Skřivan et al., 2015). One of the carotenoids suitable for this purpose is BC (Stahl and Sies, 2005), a provitamin A (Olson, 1996). In recent years, due to the increase in demand for safe animal products, it has become more important to prefer natural resources for

coloring the egg yolk (Calislar and Uygur, 2010). The nutrient profile of the eggs can be improved by diet manipulation. In the researches, it has been seen that some nutrient components that have important benefits for health can be transferred to the egg yolks via feed (Bean and Leeson, 2003; Khan et al., 2012).

Since poultry are exposed to stress conditions in a significant proportion of their lives, it is extremely important that their immune system is strong. One of the most suitable sources for this is the BC, which has a high antioxidant content. In the researches, it has been emphasized that BC enhances the survival of poultry by strengthening the immune system and positively affects the efficiency parameters.

BC is the primary source of vitamin A. Vitamin A is necessary for healthy development of bone, skin and mucosa, especially in eyesight (Thomas, 2006). Therefore, it is seen that BC has a combined effect in poultry. Particularly due to its antioxidant properties, it is thought that BC will contribute positively to the gen-

* Corresponding author email: scalislar@hotmail.com

eral immune system and performance of poultry which are exposed to diseases and effects of mycotoxins. It is also clear that as an egg component, it will have a positive impact on human health.

BC is used as a colorant and antioxidant agent in the food, cosmetics, pharmaceutical, animal feed industry (Martelli, et al., 1990; Astorg, 1997). BC gives color to products such as eggs and meat. However, it is known that most of the BC, which is used as colorant in various production industries, is synthetic. Recently, the use of natural BC has started to become widespread in line with consumer demands. By-products obtained from fruit juice production (carrot, grapefruit, apricot pulp, etc.) have low prices. Therefore, these products are used as natural BC source in chicken feeds (Sikder et. al., 1998; Mascarell et al., 2012). It has been reported that the recommended amount of BC in the feed of laying hens should be at the maximum feed rate of 30 mg / kg feed (EFSA, 2012).

It is considered that there is a small number of studies on poultry feed in relation to BC, which has important contributions to both poultry and human health, and it will be useful to focus on new research in this area.

2. Beta Carotene Resources

BC is a yellow-orange pigment. BC is found in the structure of fruits, grains, vegetables (carrots, green plants, pumpkin, spinach) and oils (Liu, 2013) with maize, green fodder, moss, marigold, stinging nettle and similar products (Table 1) (Kljak et al., 2012).

Plants such as sweet potatoes, carrots, cabbage, spinach, lettuce, fresh thyme, gourd, turnip, melon, green cabbage and broccoli are rich in BC (Groff et al., 1995). BC is more common in the leaves of plants and the amount decreases as the plant ages (Ballet et al., 2000). Egg yolks, milk, butter and liver are also animal sources containing BC. Because of its low amount and unstable structure, BC deficiency is sometimes observed in animals.

BC is also produced by algae (*Dunaliella bardawil* and *Murielopsis* sp) (Goodwin, 1992), fungi (*Blakeslea trispora*) (Mantzouridou and Tsimidou, 2008) and yeasts (*Rhodotorula glutinis*; Park et al., 2005). Among microbial sources, *Rhodotorula glutinis*, which is rich in protein, lipid and vitamins, has been reported to be suitable for producing BC (Bhosale and Gadre, 2001). In order to meet the BC needs of poultry, non-toxic *Rhodotorula* cells are used in rations (Kushwaha et al., 2014). The major carotenoid in the Western diet is BC (Stahl and Sies, 2005).

Table 1

Some sources of beta carotene and beta carotene contents

Source of BC	BC content	Literatur
Corn, yellow (mg/100g)	0.051	Yılmaz, 2010
Corn, sweet, yellow (mg/100g)	0.033	Lee et al., 1981
<i>Daucus carota</i> (mg/100 g fresh weight)	3.2-6.1	Yılmaz, 2010
Carrots, raw (mg/100g)	4.65	Bushway, 1986
Red pepper, dry (mg/100g)	2.20	Yılmaz, 2010
Peppers, sweet, red, raw (mg/100g)	0.059	Philip and Chen, 1988
Alfa alfa, green (mg/kg crude matter)	97.5	Descalzo et al., 2012
Alfa alfa, dry (mg/kg crude matter)	5.5	Descalzo et al., 2012
Soybean meal, expeller (mg/kg crude matter)	0.30	Descalzo et al., 2012
Sunflower, expeller (mg/kg crude matter)	nd	Descalzo et al., 2012
Naturel meadow grass, green (mg/kg crude matter)	63.8	Kalac, 2012
<i>Blakeslea trispora</i> (mg/L)	173	Wang, et al., 2014
<i>Rhodotorula glutinis</i> (mg/L)	6.54	Kushwaha et al., 2014

Since vitamin A cannot be synthesized by poultry, it should be taken as BC or vitamin A with feeds (Theodosiou et al., 2010). Besides carotenoids from plants, some carotenoid derivatives in the European Union have also been approved for use as an additive. These; capsantin (C40 carotenoid), P-cryptoxanthine (C40), lutein (C40), zeaxanthin (C40), P-apo-8-carotenal (C30), P-apo-8'-carotenoic acid ethyl ester (C30), xanthaxanthin (C40) and sitranaxanthin (C33) (Nimalaratne et al., 2013). The bioavailability of BC (crystal form) found in carrot juice in the feeding of wild poppies was about 30% (White et al., 1993).

Carotenoids are tetraterpenoid (C40) pigments synthesized from eight isoprene units found only in plants (Wagner and Elmadfa, 2003). They are divided into two groups according to their chemical structure: carotenes (hydrocarbon class) and xanthophylls (oxygen class) (Shete and Quadro, 2013; Von Lintig, 2012). Carotenes consist of alpha, beta and gamma carotene. The most important of carotene is the BC, which is the source of vitamin A (Taylor, 1996). BC is a fat-soluble provitamin A (Valko et al., 2007).

The BC was first isolated by Wachenroder in 1831 (Davies, 1976). The name BC was taken from carrot (*Daucus carota*) (Deming and Erdman, 1999). BC is almost always associated with chlorophyll in plants (Merck Index, 2006). The 1-carotene absorption spectrum is between 400-500 nm and is green-blue (Isler and Solms, 1971). Therefore, the BC molecule absorbs green-blue light and gives red-yellow colors.

BC is insoluble in water, acids, alkalis, but soluble in carbon disulfide and chloroform. Insoluble in meth-

anol and ethanol BC, ether, hexane and oils (FCC, 2011) slightly soluble. The diluted solution was yellow. Absorbs oxygen, which leads to inactive, colorless oxidation products (Merck Index, 2006). Pure BC is a rather dark reddish-orange color, while oxidized or melted BC is slightly yellowish orange and gray. BC, like vitamin A does not dissolve in water, it is only soluble in fat (Tek et al., 2002).

BC melts between 176-182 °C. BC, which is in the cis- and trans-isomeric forms, has a melting point of 184.50 °C (Olson, 1996). The molecular weight is 536.87 g/mol (Merck Index, 2006; FCC, 2011).

Although many carotenoids commonly have asymmetric carbon atoms, BC does not contain asymmetric carbon atoms (Woollard, 2012). The BC in the non-polar hydro-carbon group has two ion rings and theoretically this retinal structure is converted into two molecules of retinol. The conversion of carotenoids to retinol is rarely 100%. Therefore, vitamin A power of various foods is expressed as retinol equivalence (RE). Accordingly, 1 RE; 1 mg of retinol is equal to 6 mg of BC and 12 mg with other provitamin A carotenoids (Maynard et al., 1979). Vitamin A requirement of poultry is expressed as international unit (IU). It has been reported that 1 IU vitamin A activity is equivalent to 0.6 microgram BC activity or 1 mg BC is equivalent to 1.667 IU vitamin A (Blair, 2018). 1 mg of BC is equivalent to 400 IU of retinol in broiler chicks (Johannsen et al., 1998), 1200 IU in old geese and only 60 IU in young geese (Jamroz et al., 2002).

There is no proven information that the carotenoids have been transformed into another carotenoid. However, β -apo-8'-carotenal and β -apo-8' carotenoic acid ethyl esters of the BC degradation products have been shown to have coloring potential in poultry (El-Boushy and Raterink, 1992; Erdman et al., 1993). BC gives yellow-orange color to egg yolk (Dufossé, 2009).

Feed carotenoids are present in the natural compounds in about 60 to 90% trans and 10 to 30% cis form. Trans form is a more effective pigment due to its red color tone and greater stability. Chickens have the ability to convert some of the trans form of BC into the cis form and this transformation takes place in egg yolk (Hencken, 1992).

Most commercialized beta carotenes are the chemical synthesis of β -ionone (Raja et al., 2007; Ribeiro et al., 2011). The β -ionone is originally synthesized from natural sources, such as lemon grass oil or pine turpentine. However, in recent years it has been produced from β -ionone, acetone or butadiene. BC is synthesized by saponification of vitamin A acetate. Fungal and microalgae are very promising sources for the industrial production of carotenoids (Echavarri-Erasun and Johnson, 2002). Some strains of *Blakeslea trispora* fungus, a host of tropical plants, are high BC producing sources (Dufossé, 2006).

3. Functions of Beta Carotene

Provitamin A and thus BC are required to perform visual functions (Von Lintig, 2012). BC has been shown to inhibit certain types of cancer with arthrosclerosis, cataract, and multiple sclerosis due to the antioxidant properties and provitaminase activity (Terao, 1989).

BC prevents oxidative damage to cellular lipids, proteins and DNA. BC, which shows anti-inflammatory properties, protects the skin against premature aging, photodermatitis and cancers against the harmful effects of UV light (Stahl and Sies, 2007; Cazzonelli, 2011). It has been reported that carotenoids have a significant effect on skin, egg and meat quality (Liufa et al., 1997). Carotenoids have a great effect on the color of the hens' skin and egg yolk, egg and meat quality (Sirri et al., 2007; Hien et al., 2013).

The annual total carotenoid production in nature is estimated to be around 100 thousand tons. Carotenoids play an important antioxidant function by activating singlet oxygen, an oxidant formed during photosynthesis in plants (Halliwell and Gutteridge, 1999). BC is an active molecule that has properties that inactivate some reactive oxygen species in relation to its antioxidant potency. Epidemiological findings have shown that BC can prevent cancers of various organs such as lung, stomach, cervix, pancreas, colon, rectum, breast, prostate and ovary due to its antioxidant activity (Jayappriyan et al., 2013).

Carotenoids with provitamin A and antioxidant effect have cellular differentiation, growth, reproduction, gene expression, immune function, and adipocyte functions (Tourniaire et al., 2009).

According to the BC free group, cock fed with BC containing rations, has been reported to produce higher antibody titer against newcastle disease (McWhinney et al., 1989). They reported that BC used in combination with vitamin E provided more protection against the infection of *Escherichia coli* in chickens (Tengerdy et al., 1990).

According to other organs, the concentration of BC in the corpus luteum was highest but no effect on reproduction was determined (Thomas, 2006).

4. Metabolism of Beta Carotene

Vitamin A is required for the survival of all vertebrate animals. BC is one of the important sources of vitamin A requirement. Absorption of BC from intestines, transformation into vitamin A, transport, accumulation and metabolism of tissues vary according to animal species.

The conversion of BC to vitamin A generally occurs in intestinal mucosa cells and liver (Coultate, 1996). Since the BC molecule consists of a pair of retinas, two molecules of retinal formation occur when this structure is separated from the middle. However, the biological activity of BC is only about half of the retinal. The enzyme responsible for the conversion of BC to retinal is known as BC-15, 15 monooxygenase

or 15.15 si dioxigenase (Wyss et al., 2000; Dela Seña et al., 2014). Retinol and retinoic acid are also produced from the retinal (Taylor, 1996; Arıkan and Muğlalı, 1999).

Absorption of BC occurs in the duodenum of the small intestine. The absorption of BC can last for several days. Absorption is faster and more effective if there is an oil in the environment. Sometimes the BC is absorbed into the intestinal wall and is quickly converted to vitamin A in there. The rest of BC is transported in the blood as very low density lipoprotein cholesterol (Nnaji et al., 2013).

Approximately 40%-45% of total carotene content is found in egg yolk (Surai and Speake 1998; Surai et al., 1999). However, compared to other carotenoids, the amount of BC stored in the egg yolk is very low. Because BC is used as a provitamin A by poultry, it is very poor to accumulate in egg yolks or other tissues (Hammershoj et al., 2010).

Poultry predominantly accumulate oxycarotenoids in their body tissues or eggs (Goodwin 1986; Hencken 1992). The deposition rate of lutein and zeaxanthin in the egg yolk was 25%, while the accumulated amount of BC was only 0.5% (Jiang et al., 1994; Hammershoj et al., 2010).

The main storage site of BC in poultry is liver. Only 0.16% to 0.66% of the total carotenoids stored in the egg yolk of poultry cultivated under intense and semi-intensive conditions were reported as BC. It was found that the amount of BC accumulated in the duck egg yolk was 1.62% (Khan et al., 2017).

The total amount of carotenoid in the egg yolk of poultry has been reported to vary between 17.33% and 37.90%, while the amount of BC varies between 1.07% and 2.12% (Kotrbaček et al., 2013). Astaxanthin in egg yolk is stored at 14%, zeaxanthin 25% and canthaxanthin at 30-40% (Hencken, 1992).

The transfer of BC to egg yolk is 0.6% while the rate of conversion to vitamin A (5-6%) is relatively high. In a study in which chickens were given sweet potato and silage, the absorption rate of xanthophylline was 93-94% and the carotene was absorbed between 55-63% (Yamada et al., 1958). Poultry animals absorb carotenes less than xanthophylls (Surai et al., 2001). BC increase in egg yolk is only 2.1% of total carotenoids (Török et al., 2007).

The number of studies on the effects of egg yolk changes in BC content is insufficient. In previous some studies, it has been reported that the amount of BC in the egg yolk decreases due to increased storage time regardless of source (Rock et al., 1996; Thomas, 2006).

The amount of BC in the eggs of different poultry breeds varies. The amount of BC on the first day of storage white leghorn hens egg yolk has found to be 0.060 mg / g, in the first week 0.047 mg / g, in the second week 0.027 mg / g and in the third week 0.004 mg / g (Okonkwo, 2009).

5. Accumulation of Beta Carotene

Feed carotenoids can undergo numerous transformations in the metabolism of animals. Some of these compounds have vitamin A activity. Usually only monohydroxy and mono-cetocarotenoids are converted into vitamin A. Carotenoids, which have high vitamin A activity, generally have very low coloring properties (Hencken, 1992).

In a feeding study with a weight of 8000 IU vitamin A in laying hens, 80% of vitamin A was transferred to egg yolk (Squires et al., 1993). In another study, it was reported that only 85.11 micrograms of vitamin A in dietary 120 micrograms could be transferred to egg yolk (Surai et al., 1998).

The amount of BC stored in egg yolk was reported to be very low (1%) (Hammershoj et al., 2010; Xue et al., 2013). In a recent study, it was determined that 8.85% of the BC in different hybrid maize was deposited in the egg yolk of laying hens (Kristina et al., 2018).

Laying hens store vitamin A in egg yolks for incubation and embryo development during the first stages of life (Bardos, 1989). Most of the vitamin A stored in the egg yolk is retinol and a small portion is retinyl esters (Joshi et al., 1973).

Adding up to 70 g of carrots per day to the rations during the feeding of laying hens has been shown to increase the egg yolk color value, especially lutein, alpha carotene and BC content effectively (Hammershoj et al., 2010). Xanthophylls (lutein, zeaxanthin) have been found to be better absorbed than hydrocarbons carotenoids (alpha-carotene, BC) (Dumbrava et al., 2006).

The addition of lutein to the ration (100 mg / kg) increased the yolk color and redness value. Compared with the control group, lutein containing diets increased the amount of BC in egg yolks by 66%, lutein 97% and zeaxanthin by 94%. However, because it is expensive, lutein is not routinely added to rations (Englmaierová and Skininivan, 2013).

BC has an accumulation rate of less than 1% in egg yolk. It has been reported that there is a linear increase in the amount of egg yolk retinol due to the increase in the amount of BC in the diet (Jiang et al., 1994). In some previous studies, it has been reported that the amount of BC in egg yolk is 1.07-2.12 (µg / kg; Kotrbaček et al., 2013) and 0.16-1.62 (mg / kg-1; Khan et al., 2017).

Very few of the BC given with the ration passes to the yolk and the rest is converted to retinol and stored in the egg. Egg yolk color is mainly affected by fat-soluble carotenes, xanthophylls and BC. A decrease in the color of the egg yolk in line with the increase in vitamin A of the rations occurred. It has been stated that high vitamin A can cause absorption of fat-soluble pigments (Mendonça et al., 2002).

In bird species, carotenoids tend to accumulate in their immune organs. When carotenoids were included in the breeding diet, it was shown that there was a significant accumulation in the thymus and bursa fabricus of chickens. Furthermore, carotenoids from the chicken diet were still detected 4 weeks after hatching in carotenoid consuming diets fed from chickens (Koutsos et al., 2003).

Carotenoids can be exposed to oxidative effects due to storage time, room temperature and illumination (photochemical) due to the large number of double bonds in their structure. The enzymatic degradation of BC requires oxygen and the destruction at high temperatures is highest. Destruction stops after complete dehydration. Therefore, both enzymatic and photochemical effects which cause the destruction of BC during storage must be controlled (Geoffrey, 1998). Losses occur during storage of BC. It was reported that the loss in the waiting period of 25 °C for one month was 10% and the loss after three months was 29% of the initial value (EFSA, 2012).

6. Conclusion and Suggestion

Some nutrients in feeds can be transferred to eggs and functional eggs can be produced. It is thought that one of the nutrients that may contribute to functional production due to increasing the amount of egg that is passed to the egg and which is stored here may be BC. However, more information is needed about the transition of BC into eggs. It is thought that it is necessary to focus more intensely on BC, which is thought to have an important contribution to the realization of an organic and sustainable animal production suitable for human health in a century when organic egg and meat production is gaining momentum.

7. References

- Arikan Ş, Muğlalı ÖH (1999). Bazı çiftlik hayvanlarının üreme fonksiyonları üzerine β -karotenin etkisi. *Lalahan Hayvancılık Araştırma Enstitüsü Dergisi*, **39** (2): 85-94.
- Astorg P (1997). Food carotenoids and cancer prevention: An overview of current research. *Trends Food Science Technology*, **8**(12): 406-412.
- Ballet N, Robert JC, Williams PEV (2000). Vitamins in forages. In: Givens DI, Owen E, Axford RFE, Omed HM (eds) Forage evaluation in ruminant nutrition. CABI, Surrey, **19**:399-431.
- Bardos L (1989). Plasma vitamin A composition and retinol binding protein concentration during egg formation in laying hens. *International Journal for Vitamin and Nutrition Research*, **59**(3):251-254.
- Bean LD, Leeson S (2003). Long-term effects of feeding flaxseed on performance and egg fatty acid composition of brown and white hens. *Poultry Science*, **82**: 388-394.
- Bhosale P, Gadre RV (2001). β -Carotene production in sugarcane molasses by *Rhodotorula glutinis* mutant. *Journal of Industrial Microbiology and Biotechnology*, **26**(6): 327-332.
- Blair R (2018). Nutrition and feeding of organic poultry 2nd Edition. ISBN-13: 978-1786392985. 256p.
- Bushway RJ (1986). Determination of alpha- and beta-carotene in some raw fruits and vegetables by high performance liquid chromatography. *Journal of Agricultural and Food Chemistry*, **34**(3): 409-412.
- Calislar S, Uygur G (2010). Effect of dry tomato pulp on egg yolk pigmentation and some egg yield characteristics of laying hens. *Journal of Animal and Veterinary Advances*, **9**(1):96-98.
- Cazzonelli CI (2011). Carotenoids in nature: insights from plants and beyond. *Functional Plant Biology*, **38**: 833-847.
- Coulter TP (1996). Food-the chemistry of its components-2nd edit. The Royal Society of Chemistry, Cambridge.
- Davies H (1976). In: Goodwin, T.W. ed. Chemistry and biochemistry of plant pigments. London: 758 Academic Press, 1976, p 38.
- Dela Peña C, Ried KM, Narayanasamy S, Curley RWJ, Schwartz SJ, Harrison EH (2014). The human enzyme that converts dietary provitamin A carotenoids to vitamin A is a dioxygenase. *The Journal of Biological Chemistry*, **289**(19): 13661-13666.
- Deming DM, Erdman JW (1999). "Mammalian carotenoid absorption and metabolism," *Pure and Applied Chemistry*, **71**(12): 2213-2223.
- Descalzo AM, Rossetti L, Páez R, Grigioni G, García PT, Costabel L, Negri L, Antonacci L, Salado E, Bretschneider G, Gagliostro G, Comerón E, Taverna MA (2012). Differential characteristics of milk produced in grazing systems and their impact on dairy products. http://cdn.intechopen.com/pdfs/39316/InTech-Differential_characteristics_of_milk_produced_in_grazing_systems_and_their_impact_on_dairy_products.pdf Chapter 15 (Erişim tarihi: 19.03.2014).
- Dufossé L (2006). Microbial production of food grade pigments. *Food Technology and Biotechnology*, **44**(3): 313-321.
- Dufossé L (2009). Pigments, microbial. In: Schaechter, M. (Ed.), Encyclopedia of Microbiology, third ed. Elsevier/Academic Press, New York, pp. 457-471.
- Dumbrava D, Matiuti M, Druga M, Lupea A, Ianculov I, Clep C (2006). Effect of seabuckthorn berry flour from hens food on egg yolk carotenoidic pigments content. *Annals of the Faculty of Engineering Hunedoara*, **3**, 156-160.

- Echavarri-Erasun C, Johnson EA (2002). Fungal carotenoids. *Applied Mycology and Biotechnology*, **2**: 45-85.
- EFSA (2012). Scientific Opinion on the safety and efficacy of beta-carotene as a feed additive for all animal species and categories. *EFSA Journal*, **10**(6): 2737.
- El Boushy AR, Raterink R (1992). Egg yolk pigmentation. *World Review of Animal Production*, **27**(1): 49-62.
- Englmaierová M, Skřivan M (2013). Effect of synthetic carotenoids, lutein, and mustard on the performance and egg quality. In *Scientia Agriculturae Bohemica*, **44** (3): 138-143.
- Erdman JWJ, Bierer TL, Gugger ET (1993). Absorption and transport of carotenoids. *Annals of the New York Academy of Sciences*, **691**(1):76-85.
- FCC (2011). β -Carotene. In: Food Chemicals Codex. 7th ed. Rockville, MD: The US Pharma-copeial Convention. p 180. Food Chemicals Codex.
- Geoffrey LZ (1998). Biochemistry, Wm C Brown publishers, 4th edition.
- Goodwin TW (1986). Metabolism, nutrition and function of carotenoids. *Annual Review of Nutrition*, **6**, 273-297.
- Goodwin TW (1992). Distribution of carotenoids. *Methods in Enzymology*, **213**: 167-172.
- Groff JL, Gropper SS, Hunt SM (1995). Advanced nutrition and human metabolism, West Publishing Company, New York.
- Halliwell B, Gutteridge JMC (1999). Free radicals in biology and medicine. 3rd ed. New York, NY: Oxford University Press.
- Hammershoj M, Kidmose U, Steenfeldt S (2010). Deposition of carotenoids in egg yolk by short-term supplement of coloured carrot (*Daucus carota*) varieties as forage material for egg-laying hens. *Journal of the Science of Food and Agriculture*, **90**(7): 1163-1171.
- Hencken H (1992). Chemical and physiological behavior of feed carotenoids and their effects on pigmentation. *Poultry Science*, **71**(4): 711-717.
- Hien TQ, Phung TV, Tham PD, Van TT, Kien TT (2013). Feed and animal nutrition, Agri. Publishing House, Hanoi, Vietnam, 208 pp.
- Isler O, Solms G (1971). Carotenoids. Stuttgart, Germany: Birkhäuser Verlag, Basel- Stuttgart 1971. 1. Aufl., 932 S., zahlr. z. T. farbige Abb. u. Tab., geb. sfr. 118.
- Jamroz D, Jakobsen K, Wiertelcki T, Jensen SK (2002). Utilization of dietary β -carotene and retinol acetate by young and older geese. *Acta Agriculturae Scandinavica*, Section A, **52**(3): 150-158.
- Jayappriyan KR, Rajkumar R, Venkatakrishnan V, Nagaraj S, Rengasamy R (2013). In vitro anticancer activity of natural β -carotene from *Dunaliella salina* EU5891199 in PC-3 cells. *Biomedicine and Preventive Nutrition*, **3**:99-105.
- Jiang YH, McGeachin RB, Bailey CA (1994). "Alphatocopherol, beta-carotene, and retinol enrichment of chicken eggs," *Poultry Science*, **73**(7): 1137-1143.
- Johannsen AKB, Jensen SK, Jakobsen K (1998). A note on vitamin A activity of β -carotene in broilers. *Acta Agriculturae Scandinavica*, Section A. *Animal Science*, **48**(4): 260-263.
- Joshi PS, Mathur SN, Murthy SK, Ganguly J (1973). Vitamin A economy of the developing embryo and of the freshly hatched chick. *Biochemical Journal*, **136**(3): 757-761.
- Kalac P (2012). Carotenoids, ergosterol and tocopherols in fresh and preserved herbage and their transfer to bovine milk fat and adipose tissues: A review. *Journal of Agrobiology*, **29**(1): 1-13.
- Khan MS., Amin MR, Florian JS (2017). Carotenoid status of poultry egg under different feeding system in Bangladesh. *International Journal of Poultry Science*. **16**(6): 228-232.
- Khan RU, Nikousefat Z, Tufarelli V, Naz S, Javdani M, Laudadio V (2012). Garlic (*Allium sativum*) supplementation in poultry diets: Effect on production and physiology. *World's Poultry Science Journal*, **68**(3): 417-424.
- Kljak K, Drdie M, Karolyi D, Grbesa D (2012). Pigmentation efficiency of Croatian corn hybrids in egg production. *Croatian Journal of Food Technology, Biotechnology and Nutrition*, **7**, 33-37.
- Kotrbaček V, Skřivan M, Kopecký J, Pěnkava O, Hudečková P, Uhríková I, Doubek J (2013). Retention of carotenoids in egg yolks of laying hens supplemented with heterotrophic *Chlorella*. *Czech Journal of Animal Science*, **58**(5): 193-200.
- Koutsos EA, Clifford AJ, Calvert CC, Klasing KC (2003). Maternal carotenoid status modifies the incorporation of dietary carotenoids into immune tissues of growing chickens (*Gallus gallus domesticus*). *Journal of Nutrition*, **133**(4): 1132-1138.
- Kristina K, Zlatko J, Darko G (2018). Content and deposition of carotenoids in egg yolk from hens fed diets differentiated in maize hybrid. Croatian Branch of the World's Poultry Science Association, 2018. str. 334-334 (poster, međunarodna recenzija, sažetak, znanstveni).
- Kushwaha K, Saini A, Saraswat P, Agarwal MK, Saxena J (2014). Colorful world of microbes: carotenoids and their applications, Article ID 837891 Hindawi Publishing Corporation; Advances in Biology, 1-13.
- Lee CY, McCoon PE, LeBowitz JM (1981). Vitamin A value of sweet corn. *Journal of Agricultural and Food Chemistry*, **29**(6): 1294-1295.

- Liu RH (2013). Health-promoting components of fruits and vegetables in the diet. *Advances in Nutrition*, 4:384S–392S. DOI: 10.3945/an.112.003517.
- Liufa W, Xufang L, Cheng Z (1997). Carotenoids from *Alocasia* leaf meal as xanthophyll sources for broiler pigmentation. *Tropical Science*, 37(2): 116-122.
- Mantzouridou F, Tsimidou MZ (2008). Lycopene formation in *Blakeslea trispora*. Chemical aspects of a bioprocess. *Trends in Food Science and Technology*, 19(7): 363-371.
- Martelli HL, Silva DIM, Souza NO, Pomeroy D (1990). Production of β -carotene by a *Rhodotorula* strain grown on sugar cane juice. *Biotechnology Letters*, 12(3):207-208.
- Mascarell J, Carne S, Vicente A (2012). New strategies for coloring broilers with natural pigments. *All about Feed*, 20, 28-30.
- Maynard LA, Loosli JK, Hintz HF, Warner RG (1979). *Animal Nutrition*, 7th edition, pp: 13-4. McGraw Hill, New York.
- McWhinney SLR, Bailey CA, Panigrahy B (1989). Immunoenhancing effect of Beta-carotene in chicks. *Poultry Science*, 68(Suppl. 1): 94.
- Mendonça Junior CX, Almeida CRM, Mori AV, Watanabe C (2002). Effect of dietary vitamin A on egg yolk retinol and tocopherol levels. *The Journal of Applied Poultry Research*, 11(4): 373-378.
- Merck Index (2006). β -Carotene (Monograph No. 1853). In: the Merck index an encyclopedia of chemicals, drugs, and biologicals. 14th ed. Whitehouse Station, NJ: Merck and Co., Inc. p 301.
- Nimalaratne C, Wu J, Schieber A (2013). Egg yolk carotenoids: Composition, analysis, and effects of processing on their stability. In P. Winterhalter and S.E. Ebeler (Eds.), *Carotenoid cleavage products*, chapter 18: 219-225. Oxford: ACS Publications, Oxford University Press.
- Nnaji LC, Okonkwo IF, Solomon BO, Onyia OC (2013). Comparative study of beta-carotene content of egg yolk of poultry. *International Journal of Agriculture and Biosciences*, 2(1): 1-3.
- Okonkwo JC (2009). Effects of breed and storage duration on the beta-carotene content of egg yolk. *Pakistan Journal of Nutrition*. 8(10): 1629-1630.
- Olson J (1996). Carotenoid, vitamin A and cancer. *Journal of Nutrition*, 116(6): 1127-1130.
- Park PK, Cho DH, Kim EY, Chu KH 2005. Optimization of carotenoid production by *Rhodotorula glutinis* using statistical experimental design. *World Journal of Microbiology and Biotechnology*, 21(4): 429-434.
- Philip T, Chen TS (1988). Development of a method for the quantitative estimation of provitamin A carotenoids in some fruits. *Journal of Food Science*, 53(6): 1703-1706.
- Raja R, Hemaiswarya S, Rengasamy R (2007). Exploitation of *Dunaliella* for β -carotene production. *Applied Microbiology and Biotechnology*, 74(3):517-523. doi: 10.1007/s00253-006-0777-8.
- Ribeiro BD, Barreto DW, Coelho MAZ (2011). Technological aspects of β -carotene production. *Food and Bioprocess Technology*, 4(5): 693-701.
- Rock CL, Jacob RA, Bowen PE (1996). Update on the biological characteristics of the antioxidant micronutrients: Vitamin C, vitamin E and carotenoids. *Journal of the American Dietetic Association*, 96(7): 693-702.
- Shete V, Quadro L (2013). Mammalian metabolism of β -carotene: gaps in knowledge. *Nutrients*. 5(12):4849-4868. DOI: 10.3390/nu5124849.
- Sikder CA, Chowdhury DS, Rashid HM, Sikder KA, Das CS (1998). Use of dried carrot meal (DCM) in laying hen diet for Egg yolk pigmentation. *AJAS.*, 11(3): 239-244.
- Sirri F, Iaffaldano N, Minelli G, Meluzzi A, Rosato MP, Franchini A (2007). Comparative pigmentation efficiency of high dietary levels of apoester and marigold extract on quality traits of whole liquid egg of two strains of laying hens. *The Journal of Applied Poultry Research*, 16(3): 429-437.
- Skřivan M, Englmaierová M, Skřivanová E, Bubancová I (2015). Increase in lutein and zeaxanthin content in the eggs of hens fed marigold flower extract. *Czech Journal of Animal Science*, 60(3): 89-96.
- Squires MW, Naber EC (1993). Vitamin profiles of eggs as indicators of nutritional status in the laying hen: Vitamin A study. *Poultry Science*, 72(1): 154-164.
- Stahl W, Sies H (2005). Bioactivity and protective effects of natural carotenoids. *Biochimica et Biophysica Acta*, 1740(2): 101-107.
- Stahl W, Sies H (2007). Carotenoids and flavonoids contribute to nutritional protection against skin damage from sunlight. *Molecular Biotechnology*, 37(1): 26-30.
- Surai PF, Ionov IA, Kuklenko TV, Kostjuk IA, Macpherson A, Speake BK, Noble RC, Sparks NHC (1998). Effect of supplementing the hen's diet with vitamin A on the accumulation of vitamins A and E, ascorbic acid and carotenoids in the egg yolk and in the embryonic liver. *British Poultry Science*, 39(2): 257-263.
- Surai PF, McDevitt RM, Speake BK, Sparks NHC (1999). Carotenoid distribution in tissues of the laying hen depending on their dietary supplementation. *Proceedings of the Nutrition Society* 58: 30A.
- Surai PF, Speake BC, Sparks NHC (2001). Carotenoid in avian nutrition and embryonic development. 1.

- Absorption, availability and levels in plasma and egg yolk. *Journal of Poultry Science*, **38**(1): 1-27.
- Surai PF, Speake BK (1998). Distribution of carotenoids from the yolk to the tissues of the chick embryo. *Journal of Nutritional Biochemistry*. **9**(11): 645-651.
- Taylor W (1996). Response properties of long-range axon-bearing amacrine cells in the dark adapted rabbit retina. *Visual Neuroscience*, **13**(4): 599-604.
- Tek S, Kılıçarslan MR, Tek Ç, Sabuncu A (2002). Effects of beta-carotene on the fertility of rabbits, *Turkish Journal of Veterinary and Animal Sciences*, **26**(3): 497-502.
- Tengerdy RP, Lancetera NG, Nockels CF (1990). The effects of beta carotene on disease protection and humoral immunity in the chicken. *Avian Diseases*, **34**(4): 848-854.
- Terao J (1989). Antioxidant activity of beta-carotene-related carotenoids in solution. *Lipids*, **24**(7): 659-661.
- Theodosiou M, Laudet V, Schubert M (2010). From carrot to clinic: an overview of the retinoic acid signaling pathway. *Cellular and Molecular Life Sciences*, **67**(9): 1423-1445.
- Thomas MD (2006). Textbox of biochemistry with clinical correlation. John Wiley and Sons inc. 6th edn.
- Török J, Hargitai R, Hegyi G, Matus Z, Michl G, Péczely P, Rosivall B, Tóth G (2007). Carotenoids in the egg yolks of collared flycatchers (*Ficedula albicollis*) in relation to parental quality, environmental factors and laying order. *Behavioral Ecology and Sociobiology*, **61**(4): 541-550.
- Tourniaire F, Gouranton E, von Lintig J, Keijer J, Bonet ML, Amengual J, Lietz G, Landrier JF (2009). Beta-Carotene conversion products and their effects on adipose tissue. *Genes & Nutrition*, **4**(3): 179-187.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J (2007). Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry and Cell Biology*, **39**(1):44-84.
- Von Lintig J (2012). Provitamin A metabolism and functions in mammalian biology. *The American Journal of Clinical Nutrition*, **96**(suppl):1234S-1244S.
- Wagner KH, Elmadfa I (2003). Biological relevance of terpenoids. Overview focusing on mono-di- and tetraterpenes. *Annals of Nutrition and Metabolism*, **47**(3-4): 95-106.
- Wang HB, Xu RG, Yu LJ, Luo J, Zhang LW, Huang XY, Zou WA, Zhao Q, Lu MB (2014). Improved beta-carotene and lycopene production by *Blakeslea trispora* with ultrasonic treatment in submerged fermentation. *Zeitschrift für Naturforschung C*, **69**(5-6): 237-44.
- White WS, Peck KM, Ulman EA, Erdman Jr JW (1993). "The ferret as a model for evaluation of the bioavailabilities of all- trans- β -carotene and its isomers," *Journal of Nutrition*, **123**(6): 1129-1139.
- Woollard G (2012). Retinol, retinoic acid, carotenes and carotenoids: Vitamin A structure and terminology. <https://pubs.rsc.org> doi:10.1039/9781849735506-00003.
- Wyss A, Wirtz G, Woggon W, Brugge, R, Wyss M, Friedlein A, Bachmann H, Hunziker W (2000). "Cloning and expression of β , β -carotene 15,15'-dioxygenase," *Biochemical and Biophysical Research Communications*, **271**(2): 334-336.
- Xue F, Li C, Pan S (2013). In vivo antioxidant activity of carotenoid powder from tomato by product and its use as a source of carotenoids for egg laying hens. *Food & Function*, **25**(4): 610-617.
- Yamada H, Hashida A, Saito M (1958). Studies on the metabolism of carotenoids in laying hens. IV. Absorption of carotenoids in sweet potato silage and sweet potato vine silage by hens, with special reference to the formation of vitamin a in yolk. *Japanese Society of Animal Science*, **29** (2): 77-83.
- Yılmaz İ (2010). Karotenoidler. *İnönü Üniversitesi Tıp Fakültesi Dergisi*, **17**(3): 223-231.