



The Effects of Fungicidal Seed Treatments on Seed Germination, Mean Germination Time and Seedling Growth in Safflower (*Carthamus tinctorius* L.)

Nurgül ERGİN^{1,*}, Engin Gökhan KULAN², Mehmet Demir KAYA²

¹ Bilecik Şeyh Edebali University, Faculty of Agriculture and Natural Science, Department of Field Crops, Bilecik, Turkey

² Eskişehir Osmangazi University, Faculty of Agriculture, Department of Field Crops, Eskişehir, Turkey

ARTICLE INFO

Article history:

Received date: 20.02.2021

Accepted date: 29.06.2021

Keywords:

Carthamus tinctorius L.

Cultivar

Germination

Pathogen

Fungicide treatment

ABSTRACT

Seed-borne pathogens cause incorrect scores determining seed germination rate in safflower. A laboratory experiment was planned to search for an effective fungicide treatment for safflower seeds inhibiting the seed-borne infections during germination and early seedling growth. Two safflower cultivars (Olas and Linas) and five fungicides (Thiram, maneb, mancozeb, metalaxyl, and captan) were tested. Germination, mean germination time, and seedling growth parameters were investigated for determining the effectiveness of the fungicides. Results showed that germination percentage was not adversely affected by the fungicides and a higher germination percentage was obtained from the seeds treated with fungicides. Mean germination time shortened with fungicide treatments and more rapid germination was observed in thiram and metalaxyl. Fungicides increased root and shoot growth, especially in thiram and metalaxyl. The seedling weight of safflower cultivars was not changed by the application of fungicides. The infection rate of two safflower cultivars was different and untreated seeds of cv. Olas were infected. The seeds treated with fungicides successfully prevented the seed-borne infections, the minimum infection rate was obtained from the seeds treated with thiram, maneb and mancozeb. It was concluded that pretreatment of safflower seeds with thiram or mancozeb should be beneficial for avoiding seed-borne pathogens before germination test, and these applications may be tested under field conditions in terms of emergence and seed yield performance.

1. Introduction

Safflower (*Carthamus tinctorius* L.) is one of the most promising oilseed crops to meet the vegetable oil demand because it is adapted to drought and saline conditions under rainfed conditions in Turkey (Kaya et al. 2019). It belongs to the *Asteraceae* family and its seeds are called an achene. Achenes contain a single seed covered by pericarp, which makes achene suitable for placing the pathogens between seed and the pericarp. These pathogens like fungi, bacteria, and viruses adversely affect germinating seeds, seedling development, and healthy plant growth. Seed-borne pathogens prevent directly and indirectly the uniform seedling emergence and stand establishment of safflower due to reduction in germination and viability of seed (Pawar et al. 2013). For these reasons, they must be controlled by seed treatments.

Fungal pathogens are the most hazardous pathogen damaging germinating seeds and early seedling development of safflower. The most common fungi are Alternaria blight (*Alternaria carthami*), Fusarium wilt (*Fusarium oxysporum* Schlecht. f. sp. *carthami*), Verticillium wilt (*Verticillium albo-atrum*), Phytophthora root rot (*Phytophthora drechsleri*), rust (*Puccinia carthami*), brown leaf spot (*Ramularia carthami*), Cercospora leaf spot (*Cercospora carthami*), and Macrophomina root rot (*Macrophomina phaseolina*) (Chattopadhyay et al., 2015). However, several fungicides and their combinations have been extensively used for controlling these pathogens (Pawar et al. 2015). In previous studies, Sudisha et al. (2006) and Pawar et al. (2013) found a significant reduction in disease indices with pre-sowing fungicidal seed treatments. Similarly, Ellis et al. (2011) reported that seed treatments with captan and fludioxonil alleviated fungi damage by 46% and 48%, respectively over control in soybean. Adrah et al. (2020) observed that different rates of pathogen contamination on sunflower varieties and fungicide application decreased in contamination rate from 98% to less than 10%. According to Bardin et al. (2003)

* Corresponding author email: nurgulergin180@gmail.com

Pythium species, which damage seedling growth in peas, sugar beet, canola, and sunflower, decreased with Thiram application, as well as increased seedling emergence rates in the field. Inhibited seed-borne fungi in rice associated with seed treatments with mancozeb, metalaxyl, benomyl, carbendazim and thiram was reported by Ibiyam et al. (2008). In our study, it was aimed to determine if there are the beneficial effects of some fungicides for decreasing the seed-borne infection during germination and early seedling development of two safflower cultivars.

2. Materials and Methods

A laboratory experiment was conducted at the Seed Science and Technology Laboratory, Eskişehir Osmangazi University in 2020. Five fungicides, thiram (1 g kg⁻¹ seed), maneb (1.5 g kg⁻¹ seed), mancozeb (2 g kg⁻¹ seed), metalaxyl (5 g kg⁻¹ seed), and captan (2.5 g kg⁻¹ seed) were applied to the seeds of safflower cultivars Olas and Linas. Seeds without fungicidal treatment were used as control. Two hundred seeds from each cultivar were firstly weighed, put into a falcon tube and a required amount of individual fungicide was poured over the seeds. The tube was tightly closed with a cover and shaken with vortex for 5 minutes for uniform coating of seeds with the fungicides (Islam et al., 2007). Untreated seeds were used as control.

Germination test was performed by the procedures of ISTA (2003) rules with two hundred (4×50) seeds of each safflower cultivar for each fungicide. Fifty seeds were inserted into two-layer filter papers wetted with 7 ml of the distilled water for each paper. After filter papers with seeds were rolled, they were placed into a sealed plastic bag to avoid water loss. The packages were incubated at 25°C in the dark and seed with 2 mm radicle was counted every 24 h for 10 d as germinated. To evaluate the speed of germination, mean germination time (MGT) was calculated according to ISTA (2003) rules. $MGT = \frac{\sum(Dn)}{\sum n}$, where, n is the seed number germinated on day D, and D is the number of days from the beginning of the germination test. On the 10th day, ten seedlings from each treatment were randomly selected to determine the seedling growth traits such as root length (RL), shoot length (SL), seedling fresh weight (SFW), and seedling dry weight (SDW). After the seedling fresh weight was directly weighed, the seedlings were transferred into an oven at 80°C for 24 hours for determination of dry weight (Ergin et al., 2021). Also, infection rate (IR) was determined by applying the following formula:

$IR = \left(\frac{\text{Number of seeds on which infection appears}}{\text{Total number of seeds}} \right) \times 100$ (Arshad Javaid et al., 2006).

The experiment was established at two factors Completely Randomized Design (CRD) with 4 replicates. All the collected data were statically analyzed by using the MSTAT-C computer program. Means were compared with the LSD test to evaluate the differences among them (Düzgüneş et al. 1983).

3. Results and Discussion

The efficacy of fungicides on germination and seedling development is shown in Table 1. The difference between cultivars ($p < 0.01$) for mean germination time, seedling length, seedling fresh weight, seedling dry weight and infection rate were statistically significant. Differences between safflower cultivars in terms of germination percentage and root length were not significant. There were significant differences among fungicides in germination percentage, mean germination time, root length, shoot length, and infection rate. Seedling fresh and dry weights were not affected by fungicides. Two-way interaction between cultivar and fungicide was significant for mean germination time, root length, shoot length and infection rate.

Fungicide treatments for seed-borne pathogens increased the germination percentage of safflower seeds. The germination percentage ranged from 86.8% in control to 93.8% in mancozeb, shown in Table 2. Similar results were reported by Khairmar et al. (2013) and Choudhary et al. (2013) who found that germination rates in control seeds were 76-84%, while it reached up to 91-95% in the seeds with fungicide applications. Mean germination time was shortened by fungicide treatments especially thiram and metalaxyl with 1.17 days. Among the fungicides, maneb and captan gave a longer time to germinate. Sundaresh et al. (1973) and Gawade et al. (2016) determined retardation in soybean germination in untreated seeds with fungicide and inhibition due to fungal infection (46%); however, seeds treated with thiram and mancozeb were completely free from fungal growth. The seeds treated with fungicides produced longer root length compared to untreated seeds of two cultivars. metalaxyl and thiram treatments gave the longest roots with 3.88 cm and 3.51 cm, respectively. Root length of cv. Olas was more prominently increased by treatment of metalaxyl and mancozeb. Shoot length was significantly enhanced when the seeds were treated with thiram and mancozeb in cv. Olas, maneb and mancozeb in cv. Linas. In general, shoot length in thiram, maneb and mancozeb showed slightly higher or similar to control seeds. Similarly, the longest seedling length in soybean was reached in metalaxyl-M + fludioxonil application (Costa et al. 2019). Choudhary et al. (2013) reported that thiram treatment gave longer roots than captan. The results are in line with the findings of Sultana and Ghaffar (2010) who reported that fungi adversely affect seedling growth, and that healthy seedling growth can be achieved with fungicidal applications. Seedling fresh weight was not changed by fungicide treatments and similar results were observed in all treatments except for cv. Olas were treated with metalaxyl, which produced the lowest fresh weight with 183 mg plant⁻¹. Heavier fresh and dry weight was determined in cv. Linas. No significant changes were observed in seedling dry weight among the fungicides. However, Solorzano and Malvick (2011) stated that the dry weight of

corn seedlings had higher values when the seeds treated with fungicide mixtures were used.

Table 1

Analysis of variance on the investigated traits in safflower cultivars and fungicide treatments

VS	DF	Mean Square						
		GP	MGT	RL	SL	SFW	SDW	IR
Total	47	1126	2.44	32.2	7.80	4.21	0.09	22350
Cultivar (A)	1	30	0.08**	0.2	0.80**	0.64**	0.06**	2749**
Fungicide (B)	5	250*	0.35**	16.6**	2.77**	0.50	0.01	11233**
A × B	5	186	1.72**	8.3**	1.34**	0.54	0.01	5616**
Error	36	659	0.28	7.1	2.89	2.53	0.03	2751

*, **: significance level at $p < 0.05$ and $p < 0.01$, respectively. VS: Variation source, DF: Degrees of freedom, GP: Germination Percentage, MGT: Mean Germination Time, RL: Root Length, SL: Shoot Length, SFW: Seedling Fresh Weight, SDW: Seedling Dry Weight, IP: Infection Rate

Table 2

Germination percentage, mean germination time, and seedling growth parameters of two safflower cultivars treated with five fungicides

Factors	GP (%)	MGT (day)	RL (cm)	SL (cm)	SFW (mg plant ⁻¹)	SDW (mg plant ⁻¹)
<i>Cultivar</i>						
Olas	89.8	1.22 ^b	3.11 ^a	2.77 ^a	214 ^b	25.5 ^{b*}
Linan	91.3	1.30 ^a	2.98 ^b	2.51 ^b	237 ^a	32.6 ^a
<i>Fungicide</i>						
Control	86.8 ^c	1.25 ^{bc}	2.25 ^d	2.73 ^{ab}	228	29.8
Thiram	91.3 ^{ab}	1.17 ^c	3.51 ^{ab}	2.84 ^a	227	29.6
Maneb	89.3 ^{bc}	1.42 ^a	2.73 ^c	2.85 ^a	225	28.5
Mancozeb	93.8 ^a	1.24 ^{bc}	3.42 ^b	2.78 ^a	232	28.7
Metalaxyl	89.8 ^{abc}	1.17 ^c	3.88 ^a	2.19 ^c	204	28.3
Captan	92.5 ^{ab}	1.31 ^b	2.49 ^{cd}	2.45 ^c	235	29.7
<i>Cultivar × Fungicide</i>						
Olas × Control	83.5	1.10 ^{bc}	1.79 ^e	2.88 ^{bc}	230	26.7
Olas × Thiram	92.0	1.04 ^c	3.79 ^{ab}	3.30 ^a	227	26.2
Olas × Maneb	88.0	1.79 ^a	2.78 ^d	2.89 ^b	212	25.1
Olas × Mancozeb	93.0	1.16 ^{bc}	4.10 ^a	2.95 ^{ab}	224	25.1
Olas × Metalaxyl	87.0	1.09 ^{bc}	4.19 ^a	2.18 ^f	183	24.7
Olas × Captan	95.0	1.13 ^{bc}	2.04 ^e	2.42 ^{def}	207	25.3
Linan × Control	90.0	1.40 ^{abc}	2.71 ^d	2.58 ^{b-f}	227	32.9
Linan × Thiram	90.5	1.30 ^{bc}	3.23 ^{bcd}	2.38 ^{ef}	227	32.9
Linan × Maneb	90.5	1.04 ^c	2.69 ^d	2.81 ^{bcd}	238	31.9
Linan × Mancozeb	94.5	1.33 ^{bc}	2.74 ^d	2.60 ^{b-e}	241	32.2
Linan × Metalaxyl	92.5	1.24 ^{bc}	3.56 ^{abc}	2.21 ^{ef}	225	31.8
Linan × Captan	90.0	1.49 ^{ab}	2.95 ^{cd}	2.48 ^{c-f}	263	34.0

*: Means followed by the same superscript letter(s) are not significant at $p < 0.05$. GP: Germination percentage, MGT: Mean germination time, RL: Root length, SL: Shoot length, SFW: Seedling fresh weight, SDW: Seedling dry weight.

Infected seed rate is considered as the effectiveness of seed treatments by fungicides. Untreated control seeds showed that infection rate could be reached up to 100%, as in cv. Olas (Figure 1). However, fungicides inhibited the seed infection resulting from seed pathogens. The least infection rate was achieved in the application of thiram to seeds of cv. Olas. Our results confirmed the findings of Singh and Jha (2003) and Saroja (2012) who found thiram and carbendazim were the most effective agents inhibiting the growth of *Fusarium oxysporum* f.sp. *ciceri* at 1% concentration among seven fungicides. Also, in the study of Khairmar et al. (2013) was observed that the fungi in the seed my-

coflora of different safflower cultivars decreased with thiram and thiram+carbendazim applications. Suresha et al. (2012) reported that the application of carbendazim+thiram significantly reduced the seed infection compared to captan, thiram, carbendazim and carbendazim+captan applications. Moreover, the beneficial effects of fungicidal seed treatments were informed by Munkvold and O'mara (2002) in maize, Habib et al. (2007) in eggplant, Akgül et al. (2011) in peanut, Ellis et al. (2011) in soybean, Dhanamanjuri et al. (2013) in chickpea and maize, Islam et al. (2015) in wheat and Addrah et al. (2020) in sunflower.




Figure 1

Infected seed rate of safflower cultivars treated with five fungicides during germination test.

*: Values on the each column followed by same letter(s) are not significant at $p < 0.05$

In conclusion, germination test is routinely performed for seed certification processes and its results are commonly used for the required amount of seed per unit area; consequently, germination rate must be accurately determined. The main obstacle in germination tests in safflower is seed-borne pathogens which are more rapidly grown than germinating seeds. They multiply more rapidly in optimum germination conditions such as high humidity and temperature than seeds, and germination was restricted by infections. In our study, there were significant differences between safflower cultivars in terms of seed-borne pathogens. This difference results from genotypic variations because the seeds of two safflower cultivars were produced under Eskişehir conditions in 2020. But, detailed researches should be conducted by using several cultivars to make a precise decision. In addition, fungicides were effective to reduce seed-borne pathogens during the germination experiment. Pathogen-free healthy seeds are needed for desired plant populations and high yield. It is argued that seeds treatment with fungicides may favorably affect the emergence, stand establishment and consequently seed yield of safflower under field conditions. In conclusion, all the fungicides inhibited the seed-borne infections without adverse effects on germination rate and allowed to grow the healthy seedlings, and thiram, maneb and mancozeb should be advised for the seed treatments before germination test in safflower.

4. Acknowledgements

The authors thank to Ms.C. students Ş.Ö. Çetin and M.F. Kaya, and the staffs of the Seed Science and Technology Laboratory in Eskişehir Osmangazi University.

5. References

- Addrah ME, Zhang Y, Zhang J, Liu L, Zhou H, Chen W, Zhao J (2020). Fungicide treatments to control seed-borne fungi of sunflower seeds. *Pathogens* 9(1): 29.
- Akgül DS, Özgönen H, Erkilic A (2011). The effects of seed treatments with fungicides on stem rot caused by *Sclerotium rolfsii* Sacc. in peanut. *Pakistan Journal of Botany* 43(6): 2991-2996.
- Arshad Javaid AA, Akhtar N, Hanif M, Farooq MA (2006). Efficacy of some fungicides against seed-borne mycoflora of wheat. *Mycopath* 4(1): 45-49.
- Bardin SD, Huang HC, Liu L, Yanke LJ (2003). Control, by microbial seed treatment, of damping off caused by *Pythium* sp. on canola, safflower, dry pea, and sugar beet. *Canadian Journal of Plant Pathology* 25(3): 268-275.
- Chattopadhyay C, Kolte SJ, Waliyar F (2015). Diseases of edible oilseed crops. CRC Press Taylor & Francis Group. 455, New York
- Choudhary CS, Jain SC, Kumar R, Choudhary JS (2013). Efficacy of different fungicides, biocides and botanical extract seed treatment for controlling seed borne *Colletotrichum* sp. in chilli (*Capsicum annum* L.). *The Bioscan* 8(1): 123-126.
- Costa EM, Nunes BM, Ventura MV, Mortate RK, Vilarinho MS, Da Silva RM, Bessa MM (2019). Physiological effects of insecticides and fungicide, applied in the treatment of seeds, on the germination and vigor of soybean seeds. *The Journal of Agricultural Science* 11: 318-324.
- Dhanamanjuri W, Thoudam R, Dutta BK (2013). Effect of some pesticides (fungicides) on the germina-

- tion and growth of seeds/seedlings of some crop plants (i.e. *Cicer arietinum* and *Zea mays*). *Middle-East Journal of Scientific Research* 17(5): 627-632.
- Düzgüneş O, Kesici T, Gürbüz F (1983). Statistic Methods 1. Ankara Univ., Fac. of Agric. Publ. No: 861, Lesson Book No: 22, Ankara-Turkey. (in Turkish)
- Ellis ML, Broders KD, Paul PA, Dorrance AE (2011). Infection of soybean seed by *Fusarium graminearum* and effect of seed treatments on disease under controlled conditions. *Plant Disease* 95(4): 401-407.
- Ergin N, Kulan E, Gözükara M A, Kaya M F, Çetin ŞÖ, Kaya MD (2021). Response of germination and seedling development of cotton to salinity under optimal and suboptimal temperatures. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi* 24(1): 108-115.
- Gawade S, Borkar S, Suryawanshi A, Zanjare S, Shinde V (2016). Efficacy of fungicides and bioagents against internal seed borne pathogen of soybean. *Advances in Life Sciences* 28: 1250-1258.
- Habib A, Sahi ST, Ghazanfer MU, Ali S (2007). Evaluation of some fungicides against seed born mycoflora of eggplant and their comparative efficacy regarding seed germination. *International Journal of Agriculture & Biology* 9(3): 519-520.
- Ibiam OFA, Umechuruba CI, Arinze AE (2008). In vitro seed-dressing technique for the control of seed-borne fungi of rice variety Faro-29. *Journal of Applied Sciences and Environmental Management* 12(3): 39-43.
- Islam SS, Rahman MH, Hasan MJ, Ashadusjaman M, Khatun M (2007) Efficacy of fungicidal seed treatment in controlling *Alternaria* spp. in radish seed. *International Journal of Sustainable Crop Production* 2(5): 46-50.
- Islam MS, Sarker MNI, Ali MA (2015). Effect of seed borne fungi on germinating wheat seed and their treatment with chemicals. *International Journal of Natural and Social Sciences* 2(1): 28-32.
- ISTA (2003). International Rules for Seed Testing. International Seed Testing Association, Bassersdorf, Switzerland.
- Kaya MD, Akdoğan G, Kulan EG, Dağhan H, Sari A (2019). Salinity tolerance classification of sunflower (*Helianthus annuus* L.) and safflower (*Carthamus tinctorius* L.) by cluster and principal component analysis. *Applied Ecology and Environmental Research* 17(2): 3849-3857.
- Khairmar SJ, Kohire OD, Badguiar SL, Utpal D, Kamble HN (2013). Effect of fungicidal seed treatment and commercially available seed dressers on safflower seed germination and seedling vigour. *Journal of Agriculture Research and Technology* 38(1): 71-74.
- Munkvold GP, O'mara JK (2002). Laboratory and growth chamber evaluation of fungicidal seed treatments for maize seedling blight caused by *Fusarium* species. *Plant Disease* 86(2): 143-150.
- Pawar SV, Dey U, Munde VG, Sutar DS, Pal D (2013). Management of seed/soil borne diseases of safflower by chemical and biocontrol agents. *African Journal of Microbiology Research* 7(18): 1834-1837.
- Pawar K, Mishra SP, Singh RK (2015). Efficacy of bioagents and fungicides against seed borne fungi of soybean. *Annals of Plant and Soil Research* 17(1): 77-81.
- Saroja DGM (2012). Effect of fungicides on seed mycoflora and seed germination of chickpea. *Indian Journal of Plant Protection* 40(2): 150-152.
- Singh DK, Jha MM (2003). Effect of fungicidal treatment against chickpea wilt caused by *Fusarium oxysporum* f. sp. *ciceri*. *Journal of Applied Biology* 13(1/2): 41-45.
- Solorzano CD, Malvick DK (2011). Effects of fungicide seed treatments on germination, population, and yield of maize grown from seed infected with fungal pathogens. *Field Crops Research* 122(3): 173-178.
- Sudisha J, Niranjana SR, Umeha S, Prakash HS, Shetty HS (2006). Transmission of seed-borne infection of muskmelon by *Didymella bryoniae* and effect of seed treatments on disease incidence and fruit yield. *Biological Control* 37(2): 196-205.
- Sultana N, Ghaffar A (2010). Effect of fungicides, microbial antagonists and oilcakes in the control of *Fusarium solani*, the cause of seed rot, seedling and root infection of bottle gourd, bitter gourd and cucumber. *Pakistan Journal of Botany* 42(4): 2921-2934.
- Sundaresh HN, Rangnathih KG, Janardan A, Vishwanatha SR (1973). Chemical seed treatment against seed borne fungi. *Indian Phytopathology* 26: 110-111.
- Suresha V, Prasanna KPR, Pramila CK, Puttaraju TB, Balakrishna P, Siddaraja R, Surendra HS (2012). Germination and seed infection as influenced by seed treatment in vegetables. *Agricultural Science Digest* 32(4): 291-295.